

Nov. 5, 1998

Name: _____

MIDTERM TEST

1. Consider a family of square-lattice planes with $\mathbf{a}_1 = d(1, 0, 0)$ and $\mathbf{a}_2 = d(0, 1, 0)$.

- Find \mathbf{a}_3 such that this is an fcc crystal.
- Find \mathbf{a}_3 such that this is a bcc crystal.
- Find \mathbf{b}_3 for one case, indicating clearly whether you have picked the fcc or bcc direct lattice.
- Verify explicitly that this \mathbf{b}_3 gives the correct interplanar spacing (in the \hat{z} direction).

2. Consider the phonon dispersion relations $\omega_1(k) = A \sin(ka/2)$ and $\omega_2(k) = \omega_0 - B \cos(ka)$, where A and B are positive constants with $\omega_0 > A + B$, $A \gg B$, for a chain of N pairs of lighter and heavier atoms (in one dimension, with periodic boundary conditions).

- Which branch corresponds to the acoustic modes? Draw a sketch of displacements for $k \neq 0$.
 - What name is given to the other branch? Explain why.
- Determine the “critical” values ω_c of ω for which there are Van Hove singularities in the phonon density of states.
- What is the exponent α associated with this singularity (where $g(\omega) \propto |\omega_c - \omega|^\alpha$)?
- What would be required [though unphysical!] for $g(\omega)$ to go to 0 continuously in 1D?
- What is $\int \sqrt{g(\omega)} d\omega$? (Do NOT compute this explicitly; use your knowledge of what this means.) What fraction of this integral is due to the acoustic branch?
- Based on your answers to the above questions, draw a sketch of $g(\omega)$ vs. ω .

or f) At low temperature T find the power of T with which $\bullet \omega^2 @_T$ varies. (Note that this amounts to the Debye approximation at low T .)

3. In STM (scanning tunneling microscopy) experiments, one actually measures electronic density rather than atomic positions. In some materials, one sees phantom atoms: the seeming presence of a non-existent atom between two actual atoms. Is the bonding in this case primarily covalent, ionic, metallic, or van der Waals? Explain briefly.

4. Consider circles of two different radii $r^>$ and $r^<$, arranged in a checkerboard pattern.

- Solve for the critical ratio $r^>/r^<$. To what face of what ionic crystal structure is this analogous?
- Suppose the circles are replaced by two different sizes of regular [sides of equal length] octagons (oriented along the symmetry axes). Define $r^>$ and $r^<$ as the distances from the centers to the midpoint of an edge.
 - Does the critical ratio increase, decrease, or stay the same?
 - Does the

packing fraction at the critical ratio increase, decrease, or stay the same? In both cases, justify your answer.

or b') Can a regular octagon ever be the Wigner Seitz cell of a 2D Bravais lattice? Explain. (What is the largest number of sides a regular polygon can have and still be the Wigner-Seitz cell of a 2D Bravais lattice?)

5. What property of x-rays and thermal neutrons is comparable in size?

6. What two excitations are coupled to form a polariton?

7. Place the letter representing the best match in the blank in front of each of the following statements about lattice properties:

- | | |
|--|---|
| ___ Reason why in phonon dispersion $\omega(-k) = \omega(k)$. | a. Finite size of real systems |
| ___ In a scattering process, wavevector is conserved only up to a reciprocal lattice vector [$k' = k + K$] | b. Harmonic approximation of lattice vibrations |
| ___ As $k \rightarrow 0$ there is a branch with $\omega \propto k$. | c. Equipartition of energy |
| ___ Thermal conductivity does not diverge as $T \rightarrow 0$. | d. Lattice-translation invariance/ |
| ___ High-temperature lattice specific heat is independent of T. | e. Inversion symmetry |
| ___ Property of all Bravais lattices but not all lattices with bases. | f. Time-reversal symmetry |

8. In homework you used the equation $\omega^2(\mathbf{k}) = \sum_{\mathbf{R}} \sin^2\left(\frac{1}{2}\mathbf{k} \cdot \mathbf{R}\right) [A + B \frac{\mathbf{k} \cdot \mathbf{R}}{R}]$, with $A = 2\phi'(d)/d$ and $B = 2[\phi_-(d) - \phi'(d)/d]$. Verify explicitly that for a 1D Bravais chain with only nearest-neighbor interactions, this equation produces the result $\omega(k) = 2\sqrt{K/M} |\sin(ka/2)|$.