
## Phys 402 Spring 2009 Homework 7 Due Friday, April 3, 2009 @ 9 AM

## EXAM #2, covering all material up to and including this HW assignment will be held on Wednesday April 8, 2009 from 9:00 AM to 10:50 AM in Phys 1201.

- 1. Griffiths,  $2^{nd}$  Edition, Problem 9.7 2-Level Rabi flopping problem. Eliminate  $c_b(t)$  to get a  $2^{nd}$  order ODE for  $c_a(t)$ . Guess a solution of the form  $c_a(t) = Exp[\lambda \ t]$ , where  $\lambda$  is complex. Use the initial conditions on  $c_a$  and  $c_b$  to determine the unknown coefficients.
- 2. Griffiths, 2<sup>nd</sup> Edition, Problem 9.8 Stimulated vs. spontaneous emission from a blackbody radiator.
- 3. Griffiths, 2<sup>nd</sup> Edition, Problem 9.20 Magnetic Resonance a) Pauli spin matrices!
   b) Time-dependent Schrödinger equation!

Extra Credit #8
Griffiths, 2<sup>nd</sup> Edition, Problem 9.11 Selection Rules and matrix element calcualtions!



Quantum mechanics.

## Physics 402 Spring 2009 Prof. Anlage Discussion Worksheet for April 1, 2009

**1. Electric quadrupole matrix element selection rules**. Suppose we relax the constraint that the electric field is uniform over the size of an atom. By expanding the traveling wave (see Fig. 9.3) electric field  $\vec{E}(y,t) = E_0 \hat{z} \cos(ky - \omega t)$ , find the potential experienced by the electron in the atom to next order of approximation (ky <<1). This is the electric quadrupole potential. Estimate how big the correction is relative to the original term for optical radiation.

<b>2.</b> What is the form of the quadrupole matrix element? For the hydrogen atom, what selection rules on changes in the quantum number <i>m</i> arise from this type of matrix element?